The University of Burdwan
Three/Four year UG course under NEP
Paper ZOOL2011

Reptilia

GENERAL CHARACTERS AND CLASSIFICATION UPTO LIVING SUBCLASSES

DR. SAGAR ADHURYA

ASSISTANT PROFESSOR IN ZOOLOGY WEST BENGAL EDUCATION SERVICE

Introduction

Monocondyla with a scaly skin (H. Gadow. 1901).

- Reptiles are cold-blooded (ectothermic), air-breathing vertebrates characterised by dry, scaly skin and laying amniotic eggs adapted for terrestrial life. A basioccipital bone is present in the skull, which articulates with the vertebral column by a single condyle.
- Reptiles evolved from amphibian ancestors during the late Carboniferous period (~320 million years ago).
- They were among the first vertebrates to become fully terrestrial, thanks to adaptations like internal fertilisation and the amniotic egg.


L. *repere* = to crawl

General characteristics

- Habitat: Chiefly terrestrial, also found in aquatic environments (freshwater and marine).
- **Body plan:** Bilaterally symmetrical. Divisible into four regions head, neck, trunk and tail.
- **Limbs:** 2 pairs, pentadactyl. Digits with claws. Limbs are absent in a few lizards and all snakes.

• Exoskeleton: Horny epidermal scales (small), scutes (large) and dermal bony plates (osteoderms).






© Dr. Sagar Adhurya


General characteristics


- Skin: Dry and cornified, devoid of glands.
- Integumentary glands: Few scent glands, which help to attract opposite mates during breeding seasons.
- Vomeronasal organ (**Organ of Jacobson**) is well developed in most squamatas and *Sphenodons*.
- Mouth: Terminal. Jaws bear simple conical teeth. In turtle teeth is replaced by horny beaks.
- Cloacal opening: Transverse or longitudinal
- Heart: Imperfectly four-chambered. Composed of two auricles and two ventricles. Ventricles are partially divided, except, in crocodiles it is completely divided. Right and left systemic arches present.
- Blood: RBC oval and nucleated. Ectothermic



General characters

- Respiration: Mainly by the lungs. Pharyngeal and cloacal respiration is seen in many aquatic turtles.
- Kidney: Metanephric
- Excretion: Uricotelic
- Cranial nerves: 12 pairs
- Mullerian duct persists as oviduct in female and Wolffian duct is retained as vas deference in male.
- Fertilisation: Internal
- Males possess copulatory organs.







Vertebral centra

General characters

- Mostly oviparous, a few species of lizards and snakes are viviparous.
- Cleavage: Meroblastic
- Eggs: Cleidoic, i.e., large, yolky and shelled. Shells are porous, allowing gas exchange. Eggs amniotic, i.e., possess an extraembryonic membrane, amnion, chorion and allantois.
- **Development:** No metamorphosis. Youngs resemble adults. Parental care is usually absent.
- **Skull:** Single occipital condyle.
- Vertebrae: Procoelous. Two sacral vertebrae instead of single one in amphibians





Distinctive Features (Compared to Amphibians)

- **1.** Adaptation to Fully Terrestrial Life
- Reptiles are **fully terrestrial**, unlike amphibians which depend on moist environments.
- No larval stage requiring water for development.
- **1** 2. Dry, Keratinised Skin
- Skin is thick, dry, and covered with keratinised scales or scutes.
- Prevents water loss, unlike the moist, permeable skin of amphibians.


Distinctive Features (Compared to Amphibians)

- 1 3. Efficient Lungs
- Respiration occurs entirely through lungs, with no cutaneous respiration.
- Lungs are more developed and compartmentalised compared to those of amphibians.
- 4. Shelled Eggs with Extra-Embryonic Membranes
- Reptilian eggs are amniotic, with a calcareous or leathery shell.
- Contain amnion, chorion, allantois, and yolk sac enabling development on land.
- Amphibian eggs lack these and must be laid in water.

Classification of Reptilia

The class Reptilia is classified into five subclasses based on the presence/absence of certain openings through the posterolateral or temporal region of the skull.

- Anapsida: without any opening in the temporal region.
 - Found in Cotylosaurs and Chelonians
- Euryapsida: Skull with a single dorso-lateral temporal opening on either side, bounded below by postorbital and squamosal bones.
 - Found in Pelycosaurians.



supraorbital


postorbital


frontal


Fig. 1. Five types of skulls in lateral view in 5 subclasses of reptiles.


Classification of Reptilia


- Parapsida: Skull with a single dorsolateral temporal opening on either side, bounded below by the supratemporal and postfrontal bones.
 - Found in Mesosaurs and Ichthyosaurs
- **Synapsida:** Skull with a single lateral temporal opening on either side, bounded above by the postorbital and squamosal bones.
 - Found in Protorosaurs and Sauropterygians
- **Diapsida:** In this case, there are two openings on each side, separated by postorbital and squamosal bones.
 - Found in Lepidosauria and Archosauria
 © Dr. Sagar Adhurya











postorbital supratemporal fossa infratemporal fossa squamosal


DIAPSIDA

(bird stock) 10

PARAPSIDA

Fig. 1. Five types of skulls in lateral view in 5 subclasses of reptiles.

Classification (According to JZ Young (1981)



Subclass Anapsida

Characteristics

- Skull devoid of temporal fossae.
- The roof of the skull is solid.

Facts

- The most primitive form of reptile.
- The extinct order **Cotylosauria** is called **'Stem reptiles'** from which other reptiles have probably evolved.
- This subclass has only one living order: Chelonia or Testudinata.

Testudines: Tortoise, Terrapins and Turtles

- The body of the Testudines is encased by a convex dorsal shield (carapace) and a flat ventral plate (Plastron), which remains joined at the sides.
- The cloacal opening is oval or longitudinal.
- Absence of teeth in adult conditions and the jaws are covered by sharp horny plates.

Tortoise

- Shape of the shell is ovoid
- Limbs end in clawed digits, no web.
- Head and limbs can be completely withdrawn into the shell.
- Terrestrial or freshwater.



Indian star tortoise (*Geochelone elegans*)
Source: https://th.bing.com/th/id/OIP.ZRdso-HtroPWHzIVWm9CtAHaE8?rs=1&pid=ImgDe
https://th.bing.com/th/id/OIP.zrdso-HtroPWHzIVWm9CtAHaE8?rs=1&pid=ImgDe

Terrapins

- The shell is almost circular and covered with soft skin.
- Limbs are webbed, and some digits bear claws.
- Head and limbs can be completely withdrawn into the shell.
- Freshwater



Indian Flapshell Turtle (*Lissemys punctata*)
© David Raju
Source: Wikimedia Commons



Northern river terrapin (*Batagur baska*)

© Axis (ground assault 11@yahoo.com)

Taken from Northern river terrapin (Batagur baska) 15

Turtles

- The digits of limbs are ensheathed by a covering and form paddles. Limbs are clawless or clawed.
- Head cannot be withdrawn completely under the shell.
- Marine.



Olive RidleyTurtle (*Lepidochelys olivacea*)
© Curt P Finfrock

Source: iNaturalist



Loggerhead Sea Turtle (Caretta caretta)
© Roberto Pillon
Source: iNaturalist

Subclass: Lepidosauria

Gk., Lepis = scale; Squaros = Lizard

- The skull of the members of this group has two temporal vacuities, i.e., a diapsid skull.
- Lizard-like reptiles with scaled skin.

Representative of the order Rhynchocephalia



Tuatara (Sphenodon punctatus)

© Digitaltrails

Source: Flickr

- Tuatara is exclusively found in New Zealand.
- Known for a very well-developed third eye
 (Parietal organ or Pineal eye), which is made up
 of a lens, a retina with a nerve connected to the
 brain, but lacking an iris.

Lacertilia (Lizards)



Oriental Garden Lizard (Calotes versicolor)

© Devika M B

Source: iNaturalist



Yellow Monitor Lizard(Varanus flavescens)

© Sagar Adhurya

Source: iNaturalist

Ophidia (Snakes)



Indian Wolf Snake (*Lycodon aucilus*)

© Aadit Patel

Source: iNaturalist




Russel's Viper (*Daboia russeli*)

© Jonathan Hakim

Source: iNaturalist

Amphisbaenia (Worm Lizards)





Red Worm Lizard (*Amphisbaena alba*)

© Alvarovelasua

Source: iNaturalist



Black and White Worm Lizard (*Amphisbaena fuliginosa*)

© Matthieu Berroneau


Source: iNaturalist

Subclass Archosauria (Ruling reptiles)

Gk., árkhōn = leader; saúra = Lizard

- The skull is of diapsid type and lacks interparietal and parietal foramina.
- Some forms are toothless, and in others, only palatal teeth are lost.
- The lower jaw is with vacuities between dentary and angular.
- In some advanced forms, bipedality was present, and the girdles were modified accordingly.

Ruling Reptiles



Bishop PJ, Bates KT, Allen VR, Henderson DM, Randau M, Hutchinson JR. Relationships of mass properties and body proportions to locomotor habit in terrestrial © Dr. Sagar Allenyauria. Paleobiology. 2020 Nov;46(4):550-68.

Precursor of mammals

Crocodilia (Crocodiles and allies)



Saltwater Crocodile (*Crocodylus porosus*) ©Sagar Adhurya



Gharial (Gavialis gangeticus)


© Sagar Adhurya


Different crocodiles

American Alligator


Source: Animal Wildlife Blog








© Dr. Sagar Adhurya





Cuvier Dwarf Caiman

© Karel Jakubec Source: Wikimedia





Saltwater crocodile

Gharial

© kilokiilo Souce: Reddit